

# A Comprehensive Care Coordination Program's Impact on Decreasing ED Utilization

J. Moore MD, R. Modha MD, D. Bailey MHA, B. Lyon RN, A. Chess MPH, MA, C. Feher MD

Internal Medicine, Norwalk Hospital. Norwalk Community Health Centre.




1

## Introduction

13 - 27% of all US ED encounters are primary care-related ED visits estimated burden of 4.4 billion USD

Potentially preventable/avoidable with timely primary care



3

## Conclusion

High risk patient management with a dedicated case manager and comprehensive care coordinator program can significantly decrease the total number of ED admission

**References**

1. Eiland KR, Ganelin DM. Reducing preventable emergency department utilization and costs by using community health workers as patient navigators. *J Healthc Manag*. 2013;58(6):412-426.
2. Crago, Roberts, et al. "Reasons for frequent emergency department use by Medicaid enrollees: a qualitative study." *Academic Emergency Medicine* 23.4 (2016): 478-481.
3. McCord, Linda F., and Catherine W. Burt. "National hospital ambulatory medical care survey: 2003 emergency department summary." *Adv data* 358.1 (2005).
4. Pines, Jesse M., et al. "Frequent users of emergency department services: gaps in knowledge and a proposed research agenda." *Academic Emergency Medicine* 18.6 (2011): 664-669.
5. Afshar, Jorshah, et al. "Nonurgent emergency department patient characteristics and barriers to primary care." *Academic Emergency Medicine* 11.12 (2004): 1302-1310.
6. Bogley, Charles, Patrick Courtney, and Keith Burau. "Houston hospitals emergency department use study." Houston, TX(2011).
7. Cambridge Health Alliance Model of Team-Based Care Implementation Guide and Toolkit. [https://www.integration.samhsa.gov/workforce/team-members/Cambridge\\_Health\\_Alliance\\_Team-Based\\_Care\\_Toolkit.pdf](https://www.integration.samhsa.gov/workforce/team-members/Cambridge_Health_Alliance_Team-Based_Care_Toolkit.pdf)